

Two new species in the tribe *Lactuceae* (*Compositae*) from Zambia

G. V. POPE*

Summary. *Tolpis mbalensis* Pope and *Lactuca mwinilungensis* Pope are described from Zambia.

In preparation of the Flora Zambesiaca account of the tribe *Lactuceae* (*Compositae*) the following two new species need to be described.

Tolpis mbalensis Pope sp. nov. affinis *T. virgatae* Bertol. sed phyllariis paucioribus et latitudine 1 mm excedentibus, setis pappi numero plus quam 12 differt. Typus: Zambia, Richards 24540 (holotypus K).

Herba perennis; caudex lignosus, gracilis; caules solitarii vel interdum complures, usque 100 cm alti, simplices, superne ramosi, flexuosi. *Folia* caulina, plerumque 7–15·5 × 1–3·5 cm, anguste oblanceolata vel anguste obovata superiora linearis, apice acute attenuata, basi attenuata, margine remote calloso-denticulata, utrinque plus minusve sparse pilosa vel strigosa usque scabriuscula. *Inflorescentia* laxe corymboso-paniculata, pedunculis adscendentibus arcuatis. *Involucrum* usque 10 mm longum, cylindricum demum obconicum, basi truncatum vel abrupte angustatum, extra araneoso-lanatum, bracteis 2–23 subulatis subtentum; phyllaria 2-seriata, 8–10 mm longa plus minusve aequalia, anguste lanceolata, apice attenuata initio incurva, extra araneoso-glandulosa, demum reflexa. *Corolla* flava; pars inferior 5–7 mm longa, cylindrica, extra crispato-pilosa; pars superior ligulata, 7–10 mm longa, lorata, extra strigosa, lobis usque 2 mm longis extra papillatis. *Achenia* castanea, usque 3·5 mm longa, obconico-turbinata superno truncata, subtiliter 12–15-costata, glabra; pappus e setis numerosis barbellatis 5–7 mm longis constatus, setis paucis parvis interdum seriem exteriorem efficientibus.

ZAMBIA. Northern Division, Mbala (Abercorn) District; 70 km S of Mbala, 1524 m, 18 July 1930, Hutchinson & Gillett 3841 (K, SRGH); c. 3·2 km from Mbala, 1706 m, 24 April 1959, McCallum-Webster 626 (K, SRGH); Mbala, 12 July 1964, Mutimushi 854 (K, SRGH); Mwambashi Fir Plantation, 1830 m, 27 March 1952, Richards 1204 (K); Lake Chila, 1524 m, 19 May 1952, Richards 1729 (K); Lake Chila, 1650 m, 13 Feb. 1959, Richards 10855 (K); Kambole Escarpment, 1524 m, 24 April 1969, Richards 24540 (holotype K); Uningi Pans, 1524 m, 16 May 1968, Sanane 117 (K); Path to Malale Village, 1646 m, 12 Sept. 1969, Sanane 955 (K).

This species, found on the escarpments and in sandy woodlands around Mbala, is distinguished from the scapigerous and more widely distributed *T. capensis* (L.) Schultz Bip. primarily by its cauline scabridulous leaves.

Accepted for publication March 1983

*B. A. Krukoff Curator of African Botany