

omnibus minime quidem spinosis eximie differt. Suffrutex caespitosus rupicolus, ramos foliosos 40—60 cm altos apice breviter ramosos emittens. Indumentum totius plantae candidum, appresse lanato-tomentosum. Folia lanceolata acuta integerrima basi vaginiformi dilatata. Capitula in ramis terminalia. Involucrum 2·5 cm longum; phyllis 6—7-seriatis regulariter imbricatis, externis basi utrinque appendiculatis quasi lyratis, internis apice in appendicem radiato-expansum flavum nitentem dilatatis, intermediis lanceolatis acutis exappendiculatis. Receptaculum + planum paleaceum. Paleae subulatae, apice setoso-laceratae. Flores juveniles flavigeni, limbo 5-fido, a tubo vix distincto. Achaenia juvenilia dense setosa. Pappus plumosus. — Kreta: Herakleion: Insel Dia (R. 13.936).

Tragopogon lassithicus Rech. fil., n. sp. — Differt a *T. Samaritanii* Boiss. & Heldr. achaenii in rostrum per breve 1—2 mm longum crassum ab achaenio vix distinctum abeuntibus, costis acute squamoso-muricatis, involucro breviore 25 mm longo. — Kreta: Lassithi-Gebirge, Berg Lazaros, 1800—2000 m (R. 14.391).

Lactuca alpestris (Gandog.) Rech. fil., comb. nov. — Syn. *Phoenixopus alpestris* Gandog., Bull. Soc. Bot. Fr. 6: 155, 1914; Gandog., Fl. cret. 54, 1915; *Lactuca viminea* (L.) Presl var. *decumbens* H. Al., Consp. Fl. gr., Suppl. 2: 57, 1912. — Differt a *L. viminea* caulibus numerosis humilibus gracilibus, foliis parvis, imprimis caulinis minus divisis, caulinis inferioribus breviter auriculato-decurrentibus, superioribus breviter (2 mm) decurrentibus, rostro achaeniorum multo breviore, 1·5 (nec 4) mm longo. — Kreta: Psiloriti-Gebirge, 1400—2200 m (Dörfler 1100, Gandoger, R. 14.233), Lassithi-Gebirge (Gandoger, Dörfler 1061), Asterusi-Gebirge, Berg Kophinas (Baldacci II/134, Dörfler 882).

Crepis foetida L. ssp. *nova sitiaca* Rech. fil. — Proxima videtur ssp. *glandulosae* (Guss.) Fiori & Paol., a qua differt indumento longissimo densissimo e villis longis crassiulus diametrum caulis subaequantibus et e pilis brevibus tenuibus consistente ad pedunculos et ad involucrum pilis glanduliferis intermititis, caulibus numerosis humilibus validis squarroso-ramosus. — Kreta: Sitia, Südküste unterhalb Achladi (R. 12.730).

Crepis flexiscapa Rech. fil., n. sp. — Sect. *Barkhausia*. — Affinis *C. foetidae* L., a qua differt caulibus numerosis plerumque simplicibus nudis scapiformibus, foliis crassiulus glabris vel subglabris subintegris vel leviter tantum runcinato-dentatis, capitulis minutissimis, involucro florifero 6—7 mm tantum longo + 5 mm crasso, involucri phyllis inferne tantum pilis albidis brevibus glanduliferis puberulis, superne glabris, phyllis externis latioribus,