

426. Stem six or seven feet high, hollow, striated
glabrous large leaves unequally runcinate,
flowers large blue on large peduncles, down sessile.
This is perhaps a new species, it grows in
woods.

182. *Troximon pallidum* Raf. Hispidum; foliis sessiliis ovato-oblongis acutis integris ciliatis crassis,
floribus axillaribus pedonculatis adpressis, perianthis pilosis 8 partitis sub 12 floris. Raf.—
Helminia echiooides Rob. p. 426. This species
scarcely differs from *Tr. odoratum*; but it grows
in meadows, has inodorous whitish yellow flow-
ers not forming a naked raceme. Every thing
else similar.

ORDER II.—CYNAROCEPHALES.

XI. G. *ONOPIX* Raf. Perianthus ventricosus imbricatus caliculatus, caliculus parvus squamosus,
squamis perianthis apice carinatis mucronatis
plerumque spinosis. Phoranthus carnosus pilosus.
Flocculis, tubulosis inequalibus quinquefidis,
laciniis linearibus, interiora magis incisa, stigma
simplex: pappus pilosus.—Obs. This genus
which belongs to the family *Cynaridæ* having a
division of the corolla more deeply cut, differs
from *Cynara* and *Pternix* by the caliculate peri-
anthe and simple down: some species of *Chicus* or
Cirsium may perhaps be found to belong to it;
their habit is similar. The name was an ancient
one for some species of *Carduus*. Raf.