

HAB. Common in the Maturatte and other districts of the Central Province.

Folia (in meis specim.) sæpissime ad basin pinnatim 2-4-lobata.

31. NOTONIA, DC. 713. *S. Senai*

1. *N. grandiflora*, DC. Prod. vi. p. 442, cum syn.—*N. corymbosa*, DC. l. c.; Wight, Icon. t. 484.—c.p. 1743.

HAB. Batticaloa District.

32. BRACHYRAMPHUS, DC. 715. *S. Lactu*

1. *B. sonchifolius*, DC. Prod. vii. p. 177, cum syn.—*B. Heyneanus*, Wight, Icon. t. 1146. *Lactuca Heyneana*, DC. l. c. p. 140, cum syn.—c.p. 2827.

HAB. Galle, Gardner.

33. MICRORHYNCHUS, Less. 715. *S. Lactu*

1. *M. sarmentosus*, DC. Prod. vii. p. 181, cum syn.; Wight, Illust. t. 133.—c.p. 1746.

HAB. Common near the seashore.

34. SONCHUS, Linn. 715. *S. Lactu*

1. *S. cleraceus*, Linn.—c.p. 3631.

HAB. A very common weed in the Central Province, in newly-cleared ground.

35. YOUNGIA, Cass. 715. *S. Lactu*

1. *Y. lyrata*, Cass. Opusc. iii. p. 86.—*Y. Mauritiana*, *Y. Thunbergiana*, *Y. runcinata*, *Y. napifolia*, *Y. Poosia*, *Y. striata*, et *Y. ambigua*, DC. Prod. vii. pp. 192, 193, cum syn. *Y. napifolia*, Wight, Icon. t. 1147.—c.p. 1747.

HAB. Very common in the Central Province, etc., up to an elevation of 5000 feet.

2. *Y. fuscipappa*, Thw.; glabra; *caulibus* sæpe e collo plurimis, erectis, parce foliosis; *foliis radicalibus* lineari-lanceolatis, acutis, runcinatis, in petiolum longum angustatis; *caulinis* summis linearibus; *corymbis* apertis.—c.p. 1748.

HAB. Central Province, at an elevation of 5000 feet.

1-2-pedalis. *Folia* 4-8 poll. longa, $\frac{1}{2}$ - $\frac{3}{4}$ poll. lata. *Involucra* 3 lin. longa, minute bracteolata. *Achenia* fusiformia, parce compressa, puberula, striata. *Pappus* pallide fuscus.

LXXXV. STYLIDIÆ.

1. STYLIDIUM, Sw. 696

1. *S. uliginosum*, Sw. (DC. Prod. vii. p. 336.)—c.p. 3505. 3505

HAB. South of the island.