



**Fig. 1.** A portion of a strict consensus cladogram of subtribe Arecinae summarising the relationships between *Gronophyllum*, *Gulubia*, *Hydriastele* and *Siphokentia*. The cladogram results from simultaneous parsimony analysis of DNA sequence data from two low copy nuclear regions (Loo et al., in prep.). \*\* = 80 – 100% bootstrap support, \* = 60 – 79% bootstrap support.

strength of the molecular phylogenetic evidence and the morphological similarities described above, we propose a pragmatic and informative solution by reducing the four genera into a single, more broadly defined genus. In this paper, we place *Gronophyllum*, *Gulubia* and *Siphokentia* in synonymy under the oldest generic name, *Hydriastele*, and provide a synopsis of the 48 accepted species, including 34 new combinations, two new names and one new species. The species and synonymies that were recognised by the authors of the most recent accounts of the four genera (Essig 1982; Essig & Young 1985; Baker et al. 2000; Dransfield, unpublished) are accepted here. However, a revision of this group is currently in progress and it is expected that some of the 48 names will be reduced to synonymy in due course. Nevertheless, we present an outline classification here so that the new generic delimitation and validly published combinations in the genus *Hydriastele* are immediately available for general use.

### Taxonomic Treatment

**Hydriastele** H. Wendl. & Drude, Linnaea 39: 208 (1875). *Adelonenga* (Becc.) Hook. f. in Bentham & Hooker, Gen. Pl. 3 (2): 885 (1883).

**Gronophyllum** Scheff., Ann. Jard. Bot. Buitenzorg 1: 135 (1876), **synon. nov.**

*Gulubia* Becc., Ann. Jard. Bot. Buitenzorg 2: 131 (1885), **synon. nov.**

*Gulubiopsis* Becc., Bot. Jahrb. Syst. 59: 11 (1924), **synon. nov.**

*Kentia* Blume, Bull. Sci. Phys. Nat. Néerl. 1: 64 (1838), nom. illeg.

*Leptophoenix* Becc., Ann. Jard. Bot. Buitenzorg 2: 82 (1885), **synon. nov.**

*Nengella* Becc., Malesia 1: 32 (1877), **synon. nov.**

*Paragulubia* Burret, Notizbl. Bot. Gart. Berlin-Dahlem 13: 84 (1936), **synon. nov.**

*Siphokentia* Burret, Notizbl. Bot. Gart. Berlin-Dahlem 10: 198 (1927), **synon. nov.**

**1. *Hydriastele affinis* (Becc.) W. J. Baker & Loo, comb. nov.** *Nenga affinis* Becc., Malesia 1: 29 (1877). *Leptophoenix affinis* (Becc.) Becc., Ann. Jard. Bot. Buitenzorg 2: 82 (1885). *Nengella affinis* (Becc.) Burret, Notizbl. Bot. Gart. Berlin-Dahlem 13: 316 (1936). *Gronophyllum affine* (Becc.) Essig & B. E. Young, Principes 29: 136 (1985). Type: Indonesia, Papua, Kapaor, Beccari s.n. (holotype FI).

**2. *Hydriastele aprica* (B. E. Young) W. J. Baker & Loo, comb. nov.** *Gronophyllum apicum* B. E. Young, Principes 29: 139 (1985). Type: Papua New Guinea, Sandau, Telefomin, Essig & Young 74082 (holotype LAE).

**3. *Hydriastele beccariana* Burret, Repert. Spec. Nov. Regni Veg. 24: 292 (1928). Type: Indonesia, Papua, Noord R., Versteeg 1662 (holotype B†; isotypes BO, L).**

**4. *Hydriastele beguinii* (Burret) W. J. Baker & Loo, comb. nov.** *Siphokentia beguinii* Burret, Notizbl. Bot. Gart. Berlin-Dahlem 10: 198 (1927). Type: Indonesia, Maluku, Halmahera, Soa Toberoe, Beguin 1995 (holotype B†; isotype BO).

*Siphokentia pachypus* Burret, Notizbl. Bot. Gart. Berlin-Dahlem 10: 199 (1927). Type: Indonesia, Maluku, Halmahera, Weda, Beguin 2349 (holotype B†; isotype BO).

**5. *Hydriastele boumae* W. J. Baker & D. Watling sp. nov.**, *H. vitiensi* affinis sed fructu majore oblongo, triadibus decussatis dispositis et foliis indivisis in submaturitate longe persistentibus differt. Typus: Fiji, Taveuni, Bouma, Cakaudrove, Oct. 2001, Watling 170127 (holotypus K; isotypi L, NY, SUVA).

Robust, solitary tree palm to 32 m. Stem 26 – 28 cm diam., brown, nodal scars inconspicuous, adventitious root growth forming an expanded cone at base of stem