

## Palmae

by

O. BECCARI.

**Licuala** Wurmbr.

1. **Licuala spinosa** Wurmbr.  
Culi6n Isl. (MERRILL no. 543).

**Livistona** R. Br.

1. **Livistona Merrillii** Becc. — *Corypha minor* Blanco Fl. Filip. edit. 1<sup>a</sup>, p. 229; edit. Naves et Villar v. I, p. 290. — *Livistonae* sp. Naves et Villar Noviss. app. p. 281—282. — *Livistona* sp. Vidal Rev. plant. vasc. Filip. p. 279 no. 1943, 1952? — Truncus elatus. Frondium petiolus elongatus supra planiusculus, subtus obsolete angulatus, marginibus acutissimis inermibus; limbus flabellato-orbicularis in parte centrali ultra medium indivisus, 1—1,20 m diam., laciniis primariis medianis 5—6 cm latis, exterioribus angustioribus, omnibus profunde bipartitis, lobis sensim longissime acuminatis. Spadix metralis et ultra, pedunculo gracili elongato, spatha cylindracea arcte involuto, ramulis subulatis, fructibus globosis cerasi magnitudine.

Luzon Isl., Prov. Tayabas, Guinayangan (MERRILL no. 2071; January 1903, fructibus non rite maturis).

**Calamus** Linn.

1. **Calamus inflatus** Warb.  
Luzon Isl., Prov. Rizal, Bosoboso (MERRILL no. 1891); Antipolo (MERRILL no. 1641, 1643; February 1903).
2. **Calamus maximus** Blanco  
Luzon Isl., Prov. Rizal, Bosoboso (MERRILL no. 1893; April 1903).
3. **Calamus microsphaerion** Becc. n. sp. Scandens gracilis; caudices vaginati circ. 19 mm diam., vaginis inermibus; frondes cirriferæ, in parte pinnifera circ. 80 cm longæ, cirro filiformi terminatæ, petiolo brevi utrinque inermi, rachi inferne reduncaculeata, supra nuda, segmentis numerosis valde inaequidistantibus in quovis latere 2—4 irregulariter approximato-fasciculatis, tenuibus, subherbaceis, utrinque viridibus, anguste lanceolatis valde acuminatis, tenuiter 3- sub 5-costulatis, nervis