

## ATTALEA. Humb. et Kth.

1. *Attalea monosperma*. — Stipes nullus; frondibus habitu concinno; foliolis lineari-acuminatis; drupa turbinata, monosperma.

**Hab.** in variis locis provinciae Paraensis. Vulgo curuá-tinga. Fructus maturescunt in januario.

Folia 5-10 contemporanea, 5-6<sup>m</sup> longa; foliola 75<sup>m</sup>-110 utrinque, concinna, semi-oblique inserta, linearia, acuminata, lamina laterali oblique et breviter adnata. Drupa turbinata, umbonato-rostrata.

2. *A. agrestis*. — Palma humili, acauli, frondibus semicurvis; petiolo brevi, humo involuto; rachide tomentosa; foliolis lineari-lanceolatis, oblique obtusis; spadice e solo emergente, erecto; fructibus trispermis.

**Hab.** in tractibus sabulosis ad fluvium Uauinchá, confluentem amnis Jamundá. Fructus maturescunt in Augusto. Vulgo curuá-y.

(Barb. Rod. hb. Palm. 324)

Folia 5-8 contemporanea; petiolus 0<sup>m</sup>,2-0<sup>m</sup>,25 longus; rachis 1<sup>m</sup>,1; foliola oblique adnata et obtusa,