



163. *BACTRIS CIRCULARIS*, showing the annulate trunk; pinnæ, spathe and spadix.

6–7 dm. long and 4 to about 6 cm. broad, many-ribbed, acuminate, margins now and then bearing a brief very slender prickle as also the midrib underneath, rachis bearing deflexed black prickles 1–4 or 5 cm. long: spathe long, to 10 dm. and narrow, densely covered with glossy black spines 1–3 cm. long, not narrowed at base into a different manubrium or peduncle-like handle; spadix 6–8 dm. long, downwardly curved with a black-spiny peduncle, axis continuous and bearing a few stray hairs, branches 6–15 cm. long and divaricate but becoming ascending toward apex: fruit top-shaped, about 18 mm. long and broad, truncate at top with a central point, tapered to base; calyx about 4 mm. deep, strongly veined, margin closely applied and practically entire.

Tobago: in woods at Caledonia, *Broadway* 4014 (type), *Bailey* 146; Roxborough-Parlatuvier Road, *J. S. Beard* 98; near Roxborough, *Bailey* 130.

†5. *Bactris circularis*, spec. nov. Fig. 163.

Erecta ad 10 m., truncus unicus vel interdum etiam quintuplex, circa 5–10 cm. diam., circularibus albidis nudis fasciis 2–3 cm. latis et mediis liris spinarum nigrarum 2–3 cm. longarum: folia 1 m. vel plus longa; pinnæ suboppositæ et nonnullæ partes rachis vacuæ, 6–8 dm. longæ et 4–5 cm. latæ, multinervatæ subter, margines spiculis, rachis petiolusque debilibus spinis nigris 1–2 cm. longis: spatha 30 cm. longa, obsita aculeis 2–3 cm. longis; spadix laxa, saepe duplex, 25–30 cm. longa cum matura: fructus globulares, 15 mm. diam.; calyx parvus, appressus, non profunde lobatus.

Erect to 10 m., trunks four to five or sometimes one in a single stand, 5–10 cm. diam., the nodes or rings separated 2–3 cm. and bearing divaricate and deflexed black spines 2–3 cm. long (whence the name *circularis*): leaves 1 m. more or less long; pinnæ alternate or subopposite with blank spaces on the rachis, 6–8 dm. long, 4–5 cm. broad at middle of leaf, not slenderly tapered to apex, strongly many-nerved on under surface, margins rough with little points but hardly armed, rachis and petiole bearing scattered thin weak black spines 1–2 cm. long: spathe about 30 cm. long, narrow, closely covered with black prickles 2–3 cm. long, constricted toward base into a handle-like portion; spadix loose and open, perhaps double or twin, hanging in fruit and then 25–30 cm. long: fruit globular, about 15 mm. diam.; calyx small, appressed, margin shallowly lobed.

Trinidad: Melajo, St. Andrews, *Bailey* 158. A striking looking bactrid with its ridged rings of black spines and intervening whitish naked bands.

†6. *Bactris Broadwayi*, spec. nov. Figs. 164, 165.

Valde foliosa, erecta, 8–10 m. alta, trunci pauci vel unici, 5–8 cm. diam., uniformiter sed non dense aculeati: folia pendentia, 1 m. vel plus longa, æque pinnata, glauca vel albida subter; pinnæ circa 1 m. longæ, 2–3 cm. latæ, tenuiter multinervatæ, margines setiferi, rachis spinis 1–4 cm. longis: spatha 25–30 cm. longa, dense spinosa; spadix simplex, brevis, axis non continuus, rami 12–13 cm. longi: fructus pauci, pæne globosi, 4 cm. longi, 3 cm. lati, rostrum valde breve; calyx stricte appressus, in duabus continuationibus, margines obtuse lobati.