

113. CALAMUS LATIFOLIUS Roxb.—*C. inermis* T. And.
— N.-E. India.
C. LATIFOLIUS subsp. *MACRACANTHUS* Becc. — *C. macracanthus* T. And. N. E. India.
114. CALAMUS DORIAEI Becc. Burma.
115. CALAMUS MANAAN Miq. Sumatra.
116. CALAMUS SIMPLEX Becc. Malayan Peninsula.
117. CALAMUS CAESIUS Bl.—*C. glaucescens* Bl.
—Borneo; Malayan Peninsula.
118. CALAMUS OPTIMUS Becc. Borneo.
119. CALAMUS AXILLARIS Becc. Malayan Peninsula.
120. CALAMUS PALLIDULUS Becc. Malayan Peninsula.

GROUP XIV.—*Leaves* produced into a long-clawed cirrus. *Leaf-sheaths* not flagelliferous. *Primary spathes* elongate-tubular. *Fruiting perianth* (where known) not pedicelliform. *Spathellule* of fem. fl. not pedicelliform. *Seed* dorsally alveolate, not ruminant; *embryo* basilar.

A.—Scandent.

121. CALAMUS OVOIDEUS Thw. Ceylon.
122. CALAMUS ANDAMANICUS Kurz Andaman Islands.
C. ANDAMANICUS v. *NICOBARICUS* Becc. Nicobar Islands.
123. CALAMUS PALUSTRIS Griff.—*C. latifolius* (not of Roxb.)
Kurz. Mergui; Andamans; Tenasserim.
C. PALUSTRIS v. *COCHINCHINENSIS* Becc. (Pierre No. 4847).
—Cochinchina.
124. *Calamus formosanus* Becc. sp. n.—Caudicis vagina 3·35 cent. diam. apici valde gibbosa et nuda, caeterum spinis laminaribus subulatis 1-2 cent. longis armata; frondium petiolus brevissimus; rachis supra in parte basilari spinosa, subtus apicem versus aculeis reduncis simplicibus armata; segmenta valde irregulariter distributa, solitaria vel saepius in quovis latere geminatim vel ternatim remote aggregata, lanceolato-oblonga, costulis 5 nudis percursa.—*C. palustris* v. *cochinchinensi* valde affinis, differt praecipue frondium petiolo subnullo.

Formosa (Oldham in Herb. Kew).

125. *Calamus Moseleyanus* Becc. sp. n.—Frondium rachis prope basin trigona, in parte apicali subtetragona, subtus aculeis reduncis validis solitariis remote armata; segmenta non numerosa,