

except for the habit drawing on Fig. 1, which was based on a colour photograph from the type locality.

DESCRIPTION OF SPECIES

1. *Licuala bifida* Heatubun & Barfod, *spec. nov.* — Fig. 1

Palma solitaria, foliis bifidis et inflorescentia partiali spicata. — Typus: *Wally 839* (holo MAN!; iso BO, K!), Indonesia, West Papua province, Teluk Bintuni regency (previously Manokwari regency), Merdey.

Solitary, small to medium-sized palm. *Stem* to 2 m high, to 3 cm diam., internodes very short. *Leaves* c. 18 in crown; petiole 13–30 cm long, 3–6 mm wide at the base to 3–6 mm across towards the apex, adaxial face flattened at the apex and shallowly channelled at the base, abaxial face rounded, greenish yellow in colour, armed; blade small, 17–20 cm wide, simple-bifid leaves, adaxial light green, abaxial whitish, c. 112 cm long, truncate at apex and 5–10 mm wide, lobe split 30–35 cm from apex to rachis, rachis c. 40 cm long. *Inflorescences* 1 or 2 in a single palm, shorter than leaves, 82–90 cm long, branched to only 2 first-order branches; peduncle 67–75 cm long; prophyll 21–25 by 1.5–2 mm; peduncular bracts lacking; rachis elongate, c. 16 cm long; rachis bract tubular, flattened, 2-keeled, 40–50 by 1 mm, covered with brown and black indument, mouth loosely sheathing, splitting on one side, apex pointed; first-order branches close to or slightly away from mouth of rachis bract (c. 12 cm); rachillae 8–12 cm long, 2–3 mm wide. *Flowers* unknown. *Fruits* globose, 7–10 mm diam., orange when mature; seed globose 5–8 mm diam.; endocarp smooth; endosperm homogeneous.

Distribution — This has been reported from a restricted area in the Merdey valley (now Moskona Utara district of Teluk Bintuni regency), in South-Central Bird's Head Peninsula in Western New Guinea.

Habitat — This palm grows in lowland forest to 500 m asl.

Conservation status — Data deficient but probably under threat due to its restricted distribution.

Vernacular name and uses — Not recorded.

Note — The only New Guinea *Licuala* recorded so far with bifid leaves. The inflorescence structure is reminiscent of *L. graminifolia* from which it differs by normally having two partial inflorescences. Flowering material of this species has not yet been collected.

Etymology — The name refers to the entire, bifid leaves.

Additional specimen studied:

INDONESIA. West Papua: *Heatubun s.n.* (MAN!), Manokwari, Merdey district (now is Moskona Utara district of Teluk Bintuni regency), forest behind Merdey village.

2. *Licuala graminifolia* Heatubun & Barfod, *spec. nov.* — Fig. 2

Palma parva solitaria, foliis subtiliter divisus et inflorescentia non ramosa. Ab *L. bifida*, et ceteris speciebus inflorescentiam spicam praebens, folia in segmentis augustis divisus statim diagnoscenda. — Typus: *Van Royen & Sleumer 7345* (holo L!; iso K!), Indonesia, West Papua province (Vogelkop Peninsula), Kebar Valley, opposite Andjai, path to Hatai, above Aremi village, 800 m, 20 November 1961.




Fig. 2. *Licuala graminifolia* Heatubun & Barfod. a. Leaf; b. stem apex with leaf sheaths and inflorescences; c. inflorescence; d. detail of the partial inflorescence; e. flower; f. longitudinal section of flower; g. staminal ring. — Scale bar: a = 6 cm, b = 2.5 cm, c = 2 cm, d = 3 mm, e, f = 1.2 mm, g = 1 mm (all: Van Royen & Sleumer 7345, K).

Solitary, small palm. *Stem* acaulescent to about 1 m high, 1.5 cm diam., internodes short. *Leaves* c. 9 in crown, sheath disintegrating into a loose mesh of fine fibres; petiole 40–65 cm long, 2–4 mm wide at the base to 2–2.5 mm across towards the apex, adaxially flattened to shallowly channelled towards the base, abaxially rounded, unarmed; lamina fan-shaped in outline, 30–50 cm wide, divided into 9–11 segments, narrowly wedge-shaped or with slightly curved lateral margins, all segments with 1–3 adaxial ribs, more or less equal in length 20–35 cm, indentations leading to adaxial ribs 2–3 mm long, those leading to abaxial splits considerably shorter, mid-segment truncate, 6–9 mm wide at apex, basal segment obliquely truncate at apex, usually wider than the mid-segment. *Inflorescences* 1 or 2 in one palm individual, shorter than leaves, 20–60 cm long, carrying one spicate, partial inflorescence; prophyll 10–12 cm long and up to 10 mm wide, loosely fitting, splitting on one side; peduncle as measured to insertion of rachis bract 8–20 cm long; peduncular bracts lacking; rachis bract tubular, 5–9 cm long and up to 7 mm wide, covered with scattered ferruginous rammenta basally, loosely fitting to weakly inflated towards the apex, with 2 or 3 splits, 5–12 mm long; peduncle of partial inflorescence equal to or slightly exceeding rachis in length; rachilla 6–8 cm long, 1–2 mm wide. *Flowers* solitary, hermaphroditic, bullet-shaped 2.5–3 mm long; flowers borne on elevated points, subtending bract c. 1 mm long, bracteole more or less conspicuous at a lateral position, pedicel very short, calyx 2.2–2.8 mm long, hyaline distally, with three 0.4–0.6 mm long pointed lobes; corolla 2–2.2 mm long, splitting deeply to c. 5 mm above the base; androecium fused to corolla for 0.5–0.6 mm, staminal ring c. 0.5 mm high; anthers inconspicuously inserted in 2 levels, 0.5–0.6 mm long; ovary 0.7–0.8 mm long, attenuate apically; style 0.6–0.7 mm long; locules in lower half. *Fruits* globose, 8–10 mm diam., green; seed 6–8 mm diam.; endocarp smooth; endosperm homogeneous.

Distribution — Known from several collections in Western New Guinea.

Habitat — This palm is distributed in lowland rain forest ranging up to 800 m asl. It is often found in association with *Vatica rassak* Blume.

Conservation status — No immediate concern since it is quite common locally but further research is required.

Vernacular name and uses — Pupuru (Jamur dialect) is the collective term for species of *Licuala* in the Lake Jamur area. No specific use of this palm has been recorded.

Note — *Licuala graminifolia* is similar to *L. bifida* in the inflorescence structure; but in contrast to this, the inflorescence is unbranched with only one, spicate, partial inflorescence. The leaf is easily distinguished by the fine segmentation of the blade.

Etymology — The name refers to the grass-like leaf segments.

Additional specimens studied:

INDONESIA. West Papua: *Heatubun 332* (K!, MAN!), Kaimana regency (previously Fakfak regency), Teluk Etna district, km 45 logging road to Lake Jamur; *Heatubun 334* (MAN!), idem.

ACKNOWLEDGEMENTS

The authors are grateful to the keepers of the herbaria BM, BO, BRI, CAN, K, L, LAE, MAN and MO who sent material of *Licuala* on loan to AAU. The fieldwork to Lake Jamur and Sorong Selatan was funded by Royal Botanic Gardens, Kew through the Palms of New Guinea Project. The plates were skilfully rendered by Lucy T. Smith for the Palms of New Guinea Project. Dr. William Baker

kindly granted us permission to use them in this paper. CDH would like to thank the staff of AAU, K, and L for ground support and hospitality during his visits in June-July 2007.

REFERENCES

- Baker, W. & J. Dransfield. 2006. Field guide to palms of New Guinea. Royal Botanic Gardens, Kew.
- Banka, R. & A.S. Barfod. 2004. A new spectacular species of Licuala (Arecaceae, Coryphoideae) from New Guinea. Kew Bulletin 59: 73–75.
- Barfod, A.S. 2000. A new species of Licuala from New Guinea. Palms 44: 198–201.
- Barfod, A.S., R. Banka & J.L. Dowe. 2001. Field guide to palms in Papua New Guinea. AAU reports 40: 1–77.