

simpliciter ramosis, ramis longis 9—12, pendulis, floribus subrotundis, petalis, latere minime distantibus, flavovirentibus.

Plantam femineam non vidi.

Patria: N. Granada.

Caudex diametro 5—6 lin., annulis 3—4 poll. distantibus. Frondes $2\frac{1}{4}$ — $2\frac{1}{2}$ ped. longae, vagina 6—7 poll., petiolo 3 poll., rhachi 12 poll., pinnis 7—9 poll. longis, 2— $2\frac{1}{2}$ poll. latis. Spadices masculi 15 poll. longae, ramis 9 poll. longis.

Proxime affinis Ch. Lindeniana mihi, sed pinnis minoribus et floribus minus apertis sat differt.

19. Ch. gracilis Willd. in Act. Berol. 1804 p. 41. Ejusd. spec. plant. IV. 2 p. 800. Mart. Hist. II. p. 3, III. p. 160. Kunth Enum. III. p. 172 no. 6. Klotzsch in Linnaea 1847 p. 446. Borassus pinnatifrons Jeq. Hort. Schoenbr. II. 65 t. 247, 248.
20. Ch. lepidota Hrm. Wendl. in Otto et Dietr. Gartenz. 1853 no. 18 p. 138.
21. Ch. Schiedeana Mart. in Linnaea 1830 p. 204. Link in Otto et Dietr. Gartenz. 1834 no. 20 p. 153, t. 6 B. Mart. in Otto et Dietr. Gartenz. 1834 no. 31 p. 245. Ejusd. Hist. III. p. 158, t. 138, t. R. fig. II. IV. V., t. V. fig. I., t. Z. fig. III. 1—9, t. Z. II. fig. XIII., t. Z. V. fig. VI., t. Z. IX. fig. II. 1—41, t. Z. XVII. fig. VIII., t. Z. XXIII. fig. I. II. Kunthia xalappensis in Otto et Dietr. Gartenz. 1833 no. 1 p. 1.
22. ? Ch. graminifolia sp. n. Caudice remote annulato; frondibus erectis, pinnatis, vaginis cylindricis, apice paululo apertis et subauriculatis, petiolis subrotundis, basi supra callosis, evanescente canaliculatis, rhachi supra acutangula, vagina, petiolo rhachique dorso fascia dilute viridi obsolete

notatis, pinnis utrinque 36—42 approximatis, aequidistantibus, linear-lanceolatis, longissime acuminatis, horizontaliter patentibus, nervo medio nervis intramarginalibus validiore, atroviridulis; spadicibus masculis erectis, ramosis, ramis longissimis pendulis, spathis jam inter florescentiam exaridis.

Plantam femineam non vidi.

Patria: Guatemala?

Caudex diametro 9—11 lin., annulis 5—8 poll. distantibus. Frondes 5—6 ped. longae, vagina 12—16 poll., petiolo 12—15 poll., rhachi $2\frac{1}{2}$ — $2\frac{3}{4}$ ped., pinnis 11—12 poll. longis, $\frac{1}{2}$ poll. latis. Spadicum masculorum rami 12 poll. longi.

β. pinnis extimis retroflexo-patentibus.

23. Ch. elatior Mart. in Linnaea 1830 p. 205. Ejusd. Hist. III. p. 157 descr. fruct. Hrm. Wendl. in Otto et Dietr. Gartenz. 1853 no. 22 p. 171. Kunth Enum. III. p. 171 no. 1. descr. fruct.
24. Ch. desmoncoides Hrm. Wendl. in Otto et Dietr. Gartenz. 1853 no. 23 p. 177.
25. Ch. resinifera Hrm. Wendl. in Otto et Dietr. Gartenz. 1853 no. 23 p. 179.
26. Ch. scandens Liebm. in Oversigt. 1845 p. 7 I. 9. Ejusd. in litt. ex Mart. Hist. III. p. 308 no. 16. Walp. Annal. III. p. 457 no. 7.
27. Ch. affinis Liebm. in Oversigt. 1845 p. 7 I. 10. Ejusd. in litt. ex Mart. Hist. III. p. 308 no. 17. Walp. Annal. III. p. 457 no. 8.

Euchamaedoreis verisimiliter adnumeranda sunt duae species pinnis in greges approximatis:

28. Ch. Klotzschiana sp. n. Caudice erecto, annulato; frondibus erecto-patentibus, pinnatis, vaginis cylindricis apice paululo apertis, petiolis subcylindricis, basi supra callosis, subcanaliculatis, rhachi supra acutangula, vagina, petiolo rhachique dorso fascia albida obsolete notatis, pinnis utrin-