

praeditae (quorum solummodo paria 1-2 apicalia cuniculum completum formantia) adjectis annulis 1-2 minoribus, horizontaliter setigeris, inter copulatos solitariis; frondium petiolus brevis biconvexus angulis obtusis et aculeis reduncis brevibus armatus; segmenta lata, pauca (10-15) per acervulos 3-5 approximatae, in facie superiore nervis 3 vix spinulosis percursa, subtus nudae.—*D. mirabili* proximus.

Cultivated at Buitenzorg, from Palembang in Sumatra.

61. **Daemonorops Forbesii** Becc. sp. n.—Caudices vaginati 12 mill. diam.; vaginae prope apicem annulis 1-2 completis reversis, caeterum annulis minoribus incompletis prope basin interruptis et obliquis ornatae; frondium petiolus biconvexus, angulis obtusis aculeis reversis brevibus armatis; segmenta lanceolato-elongata non numerosa (27) in 3 acervulos appropinquata, in facie superiore nervis 5 setulosis percursa, subtus tantum in costa media setosa; spadix axillaris nutans subpendulus, 60 cent. longus, parte pedicellari aculeata gracili 25 cent. longa, fructibus globoso-ovoideis abrupte umbonata-mucronatis, 2 cent. longis, 18 mill. latis, semine globoso-subcompresso.

Sumatra (Forbes in Herb. Calc.)

62. **Daemonorops annulatus** Becc. sp. n.—Caudices vaginati 15 mill. diam.; vaginae annulis reversis completis late membranosis numerosis (6-7) ornatae; quorum 1-2 superioribus annulis sursum versis appropinquatis et cuniculum clausum (formicarium?) formantibus, praeterea usque ad basin annulis transversis solitariis horizontalibus praeditae; frondium petiolus biconvexus, aculeis reduncis brevibus radis ad margines obtusissimos necnon in dorso armatus; segmenta non numerosa in acervulos 3 valde remotos appropinquata, elongato-oblongata, supra 3-costulata, utrinque pilis, setis aut spinulis destituta.

N. E. Borneo; Lawas River (Burbidge in Herb. Kew).

63. **DAEMONOROPS SABUT** Becc. . . . . Malayan Peninsula.

64. **DAEMONOROPS OLIGOPHYLLUS** Becc. . . . . Malayan Peninsula.

65. **DAEMONOROPS MACROPHYLLUS** Becc. . . . . Malayan Peninsula.

66. **Daemonorops collariferus** Becc. sp. n.—Caudices vaginati 15-22 mill. diam.; vaginae breves, apice non gibbosae, ibique in ore et seriatim ad petioli basin spinis rectis pallidis longissimis (5-9 cent.) armatae, praeterea annulis setuligeris tantum reversis (4-5), quorum 2 superioribus inter eos valde appropinquatis et late membranosis, ornatae, annulis inferioribus minoribus et saepe