

11. *Daemonorops stenophyllus* Becc. sp. n.—Scandens; caudices vaginati, 2-3 cent. diam.; vaginae infra petiolum vix gibbosae, in ore nudae; caeterum undique spinis complanatis laminaribus e basi lata subulatis, majoribus 1-2 cent. longis, solitariis vel basi confluentibus et subseriatis armatae; petiolus brevissimus vel subnullus; rachis longe cirrifera in dorso aculeis solitariis prope apicem ternis et semiverticillatis armata; segmenta numerosissima aequidistantia angustissima acuminatissima, majora 35 cent. longa, 1 cent. lata, chartacea, supra in costulis 3 tenuibus remote spinulosa, subtus nuda vel costa media spinulis raris obsita, marginibus fere nudis vel remotissime et adpresse spinulosis; spadix ♀ inapertus ventricosofusiformis erectus subsessilis; spatharum rostrum nudum et inerme corpus aequans, spatha extima superficialiter bicarinata spinis pollicaribus complanato-laminaribus et e basi lata subulatis saepe pectinato-laciniatis, sparsis, vel interdum basi confluentibus et subseriatis; fructus sphaerici umbonati, 15-16 mill. diam., squamis nitidis luride stramineis, linea intramarginali vix discolore, semine globoso-subdepresso.

Sumatra; Sungei bulu near Padang (Beccari P. S. No. 909).

12. *DAEMONOROPS INTERMEDIUS* (*Calamus* Griff.) Mart.

—Malayan Peninsula.

*D. INTERMEDIUS* v. *NUDINERVIS* Becc. Singapore.

13. *DAEMONOROPS SEPAL* Becc. . . . . Malayan Peninsula.

*D. SEPAL* v. *SPHAEROCARPUS* Becc. . . . . Malayan Peninsula.

14. *DAEMONOROPS PSEUDO-SEPAL* Becc. Malay an Peninsula.

15. *Daemonorops imbellis* Becc. sp. n.—Caudex . . . ; vaginae . . . ; frondes non cirriferae (semper?) rhachi subtus rotundata, supra acute bifaciali utrinque omnino inermi; segmenta numerosa concinna aequidistantia anguste ensiformia, majora 35-38 cent. longa, 16-17 mill. lata, acuminatissima, supra in costulis 3 acutis et in nervis 2 secundariis tenuibus, subtus tantum in costa media setosospinulosa, marginibus adpresse spinulosis; spadix fructifer pendulus laxe ramosus pedunculo longiusculo (6 cent.), breviter sed argute spinoso, suffultus; fructi majusculi exacte ovoidei abrupte mucronati circiter 3 cent. longi, 2 cent. crassi, squamis nitidis cinnamomeis linea fusca marginatis, in dorso profunde sulcatis; semine globoso 18 mill. longo, 15 mill. crasso.—*D. Sepal* Becc. affinis.

District of Perak, Malayan Peninsula (Scortechini).

16. *DAEMONOROPS MARGARITAE* (Hance) . . . . . Hong-Kong.

17. *Daemonorops Pierreanus* Becc. sp. n.—Caudex erectus 15-20 mill. diam. (?); vaginae fuscescenti-furfuraceae, spinis