

11. **Daemonorops stenophyllus** Becc. sp. n.—Scandens; caudices vaginati, 2-3 cent. diam.; vaginae infra petiolum vix gibbosae, in ore nudae; caeterum undique spinis complanatis laminaribus e basi lata subulatis, majoribus 1-2 cent. longis, solitariis vel basi confluentibus et subseriatis armatae; petiolus brevissimus vel subnnullus; rachis longe cirrifera in dorso aculeis solitariis prope apicem ternis et semiverticillatis armata; segmenta numerosissima aequidistantia angustissima acuminatissima, majora 35 cent. longa, 1 cent. lata, chartacea, supra in costulis 3 tenuibus remote spinulosa, subtus nuda vel costa media spinulis raris obsita, marginibus fere nudis vel remotissime et adpresso spinulosis; spadix ♀ inapertus ventricoso-fusiformis erectus subsessilis; spatharum rostrum nudum et inerme corpus aequans, spatha extima superficialiter bicarinata spinis pollicariibus complanato-laminaribus et e basi lata subulatis saepe pectinato-laciiniatis, sparsis, vel interdum basi confluentibus et subseriatis; fructus sphaerici umbonati, 15-16 mill. diam., squamis nitidis luride stramineis, linea intramarginali vix discolore, semine globoso-subdepresso.

Sumatra; Sungai bulu near Padang (Beccari P. S. No. 909).

12. **DAEMONOROPS INTERMEDIUS** (*Calamus* Griff.) Mart.

—Malayan Peninsula.

D. INTERMEDIUS v. NUDINERVIS Becc. . . . Singapore.

13. **DAEMONOROPS SEPAL** Becc. . . . Malayan Peninsula.

D. SEPAL v. SPAHEROCARPUS Becc. . . . Malayan Peninsula.

14. **DAEMONOROPS PSEUDO-SEPAL** Becc. . . . Malayan Peninsula.

15. **Daemonorops imbellis** Becc. sp. n.—Caudex . . . ; vaginae . . . ; frondes non cirriferae (semper?) rhachi subtus rotunda, supra acute bifaciali utrinque omnino inermi; segmenta numerosa concinna aequidistantia anguste ensiformia, majora 35-38 cent. longa, 16-17 mill. lata, acuminatissima, supra in costulis 3 acutis et in nervis 2 secundariis tenuibus, subtus tantum in costa media setoso-spinulosa, marginibus adpresso spinulosis; spadix fructifer pendulus laxe ramosus pedunculo longiusculo (6 cent.), breviter sed argute spinoso, suffultus; fructi majusculi exakte ovoidei abrupte mucronati circiter 3 cent. longi, 2 cent. crassi, squamis nitidis cinnamomeis linea fusca marginatis, in dorso profunde sulcatis; semine globoso 18 mill. longo, 15 mill. crasso.—*D. Sepal* Becc. affinis.

District of Perak, Malayan Peninsula (Scortechini).

16. **DAEMONOROPS MARGARITHAE** (Hance) . . . Hong-Kong.

17. **Daemonorops Pierreanus** Becc. sp. n.—Caudex erectus 15-20 mill. diam. (?); vaginae fuscescenti-furfuraceae, spinis