

dium spadiceis vix superante, exteriore interiore duplo brevior; fl. ♂ corolla calyce duplo longiore.

*Hab.*—In sylvis nunquam inundatis ad fl. Javary; floret mense Decembro. “Tamanduá pecu,” vel “Tamanduá ubim” nuncupatur.

2. *G. acaulis*, Mart., subsp. *Tapajotensis* (Trail hb. Palm, 9).—*G. acauli* minor; foliis brevioribus (vix 1<sup>m</sup>) et tenerioribus, pinnis 3-4-jugis rhomboideo-lanceolatis, .22<sup>m</sup>-.35<sup>m</sup> × .035<sup>m</sup>-.09<sup>m</sup>; venis primariis utrinque 20-22, bis flexis; angulum costale 30°-60° medio efformantibus; petiolo (vagina inclusa) .45<sup>m</sup>-.55<sup>m</sup> supra canaliculato; spadicebus .5<sup>m</sup>-.6<sup>m</sup>, rachibus .05<sup>m</sup>-.075<sup>m</sup> × .003<sup>m</sup> cauda sterili .012<sup>m</sup> mucronatis; alveolis 6-7-stichis; spathis coriaceo-membranaceis, interiore .14<sup>m</sup>-.15<sup>m</sup>, exteriore .05<sup>m</sup>-.06<sup>m</sup>; floribus *G. acaulis*; fructibus (vix maturis) globosis, pisi minoris magnitudine.

*Hab.*—In sylvis humidis, hiberno inundatis, ad Arama-hy prope fl. Tapajoz; flores et fructus inveni mense Januario.

3. *G. CAMANÁ*, sp. n., an *G. Jussieuana*, Mart., subsp. (Trail hb. Palm, 182).—Caudescens, caudice 1.0<sup>m</sup>-1.4<sup>m</sup> × .025<sup>m</sup>-.040<sup>m</sup> erecto vel flexuoso, annulis multis cincto, ligno albo, mollissimo; foliis 1.4<sup>m</sup>-2.4<sup>m</sup> subregulariter innatofissis, pinnis 17-31 (plerumque 17-21) jugis, uni- vel biveniis, angustis, subfalcato-acuminatis, apicalibus latioribus, glauco-viridibus; venis primariis utrinque 25-36, bis flexis; petiolo .6<sup>m</sup>-1.2<sup>m</sup> supra sulcato; spadicebus .45<sup>m</sup>-.60<sup>m</sup>, rachibus .15<sup>m</sup>-.20<sup>m</sup>, obtusis; alveolis 15-stichis, labio inferiore prominulo, emarginato; spathis æquilongis .225-.300<sup>m</sup>, linearibus, ancipitibus; fl. ♂ sepala petalis majora; fl. ♀ —; fructibus ovalibus, scabris nigris, .009<sup>m</sup> × .006<sup>m</sup>.

*Hab.*—In sylvis humidis ad ripas fluminum Javary et Jutahi, ab indigenis Juriti-ubim vel Assai rana (i.e., *Euterpe* sp. spuria) nuncupatur.

*Obs.*—From *G. Jussieuana*, Mart. (Palm. Orbign., p. 24 t. xv., f. 1, et t. xxiii., f. A).—This Palm differs in the longer leaves, more numerous pinnæ, blunt rachis only half as long as the peduncle, and in the oval fruits. The localities are also very distinct, *G. Jussieuana*, Mart., occurring on rocky mountain slopes, while *G. Camaná* occurs in low-lying, rather swampy forests.

\*\* Tubus stamineus ore breviter 6-dentatus.

4. *G. elegans*, Mart., var. *Amazonica* (Trail hb. Palm, 133).—Hæc varietas ab exemplis typicis in provinciis Bahiensi et Rio de Janeiro lectis differt ut infra; foliis simplicibus, bifurcis, dein varie et irregulariter pinnatisectis, laciniis utrinque 2-5, sæpe 3-jugis oppositis; venis primariis utrinque 24-28; spadicebus .10<sup>m</sup>-.30<sup>m</sup>, rachibus .05<sup>m</sup>-.20<sup>m</sup>; spathis pedunculum superantibus, laceris; alveolis 6-8-stichis; fructibus subglobosis.