

Palmer River, *Brass* 7201 (isotype A!).

17. **Gronophyllum pleurocarpum** (Burret) Essig & Young **comb. nov.** *Nengella pleurocarpa* Burret in Notizbl. Bot. Gart. Berlin-Dahlem 13: 314. 1936.

*Nengella calophylla* var. *montana* Beccari in Bot. Jahrb. Syst. 52: 27. 1914. Type: New Guinea, northeastern Papua New Guinea, Madang area, *Schlechter* 16291 (holotype B).

18. **Gronophyllum procerum** (Blume) H.E. Moore in Gentes Herb. 9: 265. 1963.

*Kentia procera* Blume, Rumphia 2: t. 106. 1838–39; 94. 1843. Type: New Guinea, southwestern West Irian, *Zippelius* s.n. (holotype L).

19. **Gronophyllum ramsayi** (Beccari) H.E. Moore in Gentes Herb. 9: 265. 1963.

*Gulubia ramsayi* Beccari in Webbia 3: 159. 1910. Type: Australia, Northern Territory, Port Essington, *Ramsay* s.n. (holotype ?MEL).

*Kentia ramsayi* (Beccari) Beccari in Webbia 4: 148. 1913.

20. **Gronophyllum rhopalocarpum** (Beccari) Essig & Young **comb. nov.**

*Nengella rhopalocarpa* (Beccari) Burret in Notizbl. Bot. Gart. Berlin-Dahlem 13: 314. 1936; *Nengella calophylla* var. *rhopalocarpa* Beccari in Bot. Jahrb. Syst. 52: 28. 1914. Type: New Guinea, Northeast Papua New Guinea, Waria River, *Schlechter* 17466 (holotype B).

21. **Gronophyllum sarasinorum** Burret in Notizbl. Bot. Gart. Berlin-Dahlem 13: 202. 1936. Type: Indonesia, Sulawesi, Posso Lake, *Sarasin* 896 (holotype B).

22. **Gronophyllum selebicum** (Beccari)

Beccari in Ann. Jard. Bot. Buitenzorg 2: 82. 1885.

*Nenga selebicum* Beccari, Malesia 1: 30. 1877. Type: Indonesia, Sulawesi, Kandari, *Beccari* s.n. (holotype FI).

Dubious species:

The following species are poorly known and considered dubious in Young's dissertation. They are therefore not included in the key to the species. A new epithet for *Nengella mayrii* is however necessary to avoid duplication with *Gronophyllum mayrii*.

**Gronophyllum affine** (Beccari) Essig & Young **comb. nov.**

*Nenga affinis* Beccari, Malesia 1: 29. 1877. Type: New Guinea, northwestern West Irian, Kapaor, *Beccari* s.n., under accession numbers 11218–11218A at FI (holotype FI).

*Leptophoenix affinis* (Beccari) Beccari in Ann. Jard. Bot. Buitenzorg 2: 82. 1885.

*Nengella affinis* (Beccari) Burret in Notizbl. Bot. Gart. Berlin-Dahlem 13: 316. 1936.

**Gronophyllum cyclopensis** Essig & Young **nom. nov.**

*Leptophoenix mayrii* Burret in Notizbl. Bot. Gart. Berlin-Dahlem 11: 709. 1933. Type: New Guinea, northwestern West Irian, *Mayr* 24 (holotype B).

*Nengella mayrii* (Burret) Burret in Notizbl. Bot. Gart. Berlin-Dahlem 13: 314. 1936.

**Gronophyllum micranthum** (Burret) Essig & Young, **comb. nov.**

*Leptophoenix micrantha* Burret in Notizbl. Bot. Gart. Berlin-Dahlem 11: 710. 1933. Type: New Guinea, West Irian, Wandammen Mtns. *Mayr* 253 (holotype B).

*Nengella micrantha* (Burret) Burret in Notizbl. Bot. Gart. Berlin-Dahlem 13: 314. 1936.